Author Index

Ackermans, M. T., Everaerts, F. M. and Beckers, J. L.
Relationship between zone length and step height in isotachophoresis 595(1992)327

Alexander, P. W., see Głod, B. K. 595(1992)149

Andersson, P. E., Demirbüker, M. and Blomberg, L. G.
Quantitative hydrocarbon group analysis of gasoline and diesel fuel by supercritical fluid chromatography 595(1992)301

Araki, T., see Kimata, K. 595(1992)77

Baumes, R. L., see Voirin, S. G. 595(1992)269

Bayonove, C. L., see Voirin, S. G. 595(1992)269

Beckers, J. L., see Ackermans, M. T. 595(1992)327

Bemgård, A., Colmsjö, A. and Lundmark, B.-O.
Gas chromatographic analysis of high-molecular-weight polynuclear aromatic hydrocarbons. I. Molecular weight 328 595(1992)247

Blaas, W., see Jiao, Y. 595(1992)364

Blomberg, L. G., see Andersson, P. E. 595(1992)301

Boutin-Muma, B., see Cohen, H. 595(1992)143

Bremner, J. B., see Richardson, D. E. 595(1992)155

Bringmann, G., Hesselmann, C. and Feineis, D.
Endogenous alkaloids in man. XI. Analysis of glyoxylate-derived 1,3-thiazolidines by ion-pair-assisted reversed-phase chromatography 595(1992)351

Brown, Jr., F. and Heath, P. C.
Separation and purification of azetidinyl methyl sulfinates using preparative high-performance liquid chromatography 595(1992)355

Camamo, M., see Haken, J. K. 595(1992)283

Chen, S.-H., Wu, S.-M. and Wu, H.-L.
Stereochemical analysis of betamethasone and dexamethasone by derivatization and high-performance liquid chromatography 595(1992)203

Clark, D., see Tharakan, J. 595(1992)103

Cohen, H. and Boutin-Muma, B.
Fluorescence detection of trichothecene mycotoxins as coumarin-3-carbonyl chloride derivatives by high-performance liquid chromatography 595(1992)143

Colmsjö, A., see Bemgård, A. 595(1992)247

Craft, N. E., see Epler, K. S. 595(1992)89

Demirbüker, M., see Andersson, P. E. 595(1992)301

Drohan, W., see Tharakan, J. 595(1992)103

Ďulák, K., see Petro, M. 595(1992)337

Dunn, M. J. and Hahn, D. A.
Aqueous high-performance size-exclusion chromatographic assay for high-molecular-weight impurities in ceftiofur sodium 595(1992)185

Epler, K. S., Sander, L. C., Ziegler, R. G., Wise, S. A. and Craft, N. E.
Evaluation of reversed-phase liquid chromatographic columns for recovery and selectivity of selected carotenoids 595(1992)89

Everaerts, F. M., see Ackermans, M. T. 595(1992)327

Feineis, D., see Bringmann, G. 595(1992)351

Francotte, E. and Wolf, R. M.
Chromatographic resolution on methylbenzoylcellulose beads. Modulation of the chiral recognition by variation of the position of the methyl group on the aromatic ring 595(1992)63

Fukuda, M., see Koizumi, K. 595(1992)340

Gerig, T. M., see Jones, L. A. 595(1992)221

Glennon, J. J., see Jones, L. A. 595(1992)209

Glennon, J. J., see Jones, L. A. 595(1992)221

Głod, B. K., Haddad, P. R. and Alexander, P. W.
Ion-exclusion chromatography using mobile phases containing β-cyclodextrin 595(1992)149

Grüner, B., Plzák, Z. and Vinš, I.
Purity assay of sodium mercaptododecaborate by high-performance liquid chromatography 595(1992)169

Gumprecht, D. L.
Comparison of commercially available thin-layer chromatography plates with mixtures of dyes, analgesics and phenols 595(1992)368

Haddad, P. R., see Głod, B. K. 595(1992)149

Hahn, D. A., see Dunn, M. J. 595(1992)185

Haken, J. K. and Camamo, M.
Degradation of polysulphones by alkali fusion 595(1992)283

Heath, P. C., see Brown, Jr., F. 595(1992)355

Hesselmann, C., see Bringmann, G. 595(1992)351

Highsmith, F., see Tharakan, J. 595(1992)103

Hlavačková, E. and Reya, R.
Comparison of colorimetric and chromatographic determination of *tert.*-butylpyrocatechol 595(1992)137

Hochuli, E., see Loetscher, P. 595(1992)113

Horie, M., see Saito, K. 595(1992)163

Hosoya, K., see Kimata, K. 595(1992)77

Itoh, H., see Kawasaki, S. 595(1992)193

Jang, M. D., see Park, J. H. 595(1992)45

Jiao, Y., Blaas, W., Rühl, C. and Weber, R.
Identification of ochratoxin A in food samples by chemical derivatization and gas chromatography–mass spectrometry 595(1992)364

Jones, L. A., Glennon, J. J. and Reiss, W. H.
Critical capillary column examination of the relationship between separation number and height equivalent to a theoretical plate and their dependence on temperature 595(1992)209

Jones, L. A., Reiss, W. H., Glennon, J. J. and Gerig, T. M.
Van Deemter-type relationship for determining the optimum initial flow-rate and optimum pressure programming rate in temperature/pressure-programmed capillary column gas chromatography utilizing separation numbers 595(1992)221

Karaiskakis, G., see Koliadima, A. 595(1992)237

Katsanos, N. A., see Koliadima, A. 595(1992)237

Kawasaki, S., Ueda, H., Itoh, H. and Tadano, J.
Screening of organophosphorus pesticides using liquid chromatography–atmospheric pressure chemical ionization mass spectrometry 595(1992)193
Kenndler, E. and Schwer, C.
Peak dispersion and separation efficiency in high-performance zone electrophoresis with gel-filled capillaries 595(1992)313
Kimata, K., Hosoya, K., Tanaka, N., Araki, T. and Patterson, Jr., D. G.
Preparation of nitrophenylethylsilylated silica gel and its chromatographic properties in the separation of polychlorinated dibenzo-*p*-dioxins 595(1992)77
Klimeš, J., Klimešová, V. and Waisser, K.
Chromatographic behaviour of dipyridylsulphides. Relationship between log k' values and structure by reversed-phase high-performance liquid chromatography 595(1992)334
Klimešová, V., see Klimeš, J. 595(1992)334
Koizumi, K., Kubota, Y., Ozaki, H., Shigenobu, K., Fukuda, M. and Tanimoto, T.
Analyses of isomeric mono-O-methyl-D-glucoses, D-glucobioses and D-glucose monophosphates by high-performance anion-exchange chromatography with pulsed amperometric detection 595(1992)340
Koliadima, A., Karaiskakis, G., Katsanos, N. A. and Roth, M.
Activity coefficients in binary liquid mixtures measured by reversed-flow gas chromatography 595(1992)237
Križanovič, K., see Petro, M. 595(1992)337
Kubota, Y., see Koizumi, K. 595(1992)340
Kulik, W., see Van Rhijn, J. A. 595(1992)289
Lagomarsino, R. J. and Latner, N.
Wide dynamic range electron-capture detection using the electron-capture detector pulse frequency signal 595(1992)359
Lang, V.
Polychlorinated biphenyls in the environment 595(1992)1
Latner, N., see Lagomarsino, R. J. 595(1992)359
Lee, T. T. and Yeung, E. S.
High-sensitivity laser-induced fluorescence detection of native proteins in capillary electrophoresis 595(1992)319
Lindner, W.
Chiral separations by liquid chromatography (edited by S. Ahuja) (Book Review) 595(1992)375
Loetscher, P., Mottlau, L. and Hochuli, E.
Immobilization of monoclonal antibodies for affinity chromatography using a chelating peptide 595(1992)113
Lundmark, B.-O., see Bemgård, A. 595(1992)247
Monger, L. S. and Olliff, C. J.
Reversed-phase chromatographic behaviour of β-endorphin: evidence of conformational change 595(1992)125
Mottlau, L., see Loetscher, P. 595(1992)113
Müller, H.-M., see Schwadorf, K. 595(1992)259
Nagae, N., see Takeuchi, T. 595(1992)121
Nakagomi, K., see Saito, K. 595(1992)163
Nakazawa, H., see Saito, K. 595(1992)163
Nose, N., see Saito, K. 595(1992)163
O'Grady, B. V., see Richardson, D. E. 595(1992)155
Olliff, C. J., see Monger, L. S. 595(1992)125
Ozaki, H., see Koizumi, K. 595(1992)340
Park, J. H., Jang, M. D. and Shin, M. J.
Solvatochromic hydrogen bond donor acidity of cyclodextrins and reversed-phase liquid chromatographic retention of small molecules on a β-cyclodextrin-bonded silica stationary phase 595(1992)45
Patterson, Jr., D. G., see Kimata, K. 595(1992)77
Petro, M., Križanovič, K. and Ďulák, K.
High-performance liquid chromatography of isopropylphenols 595(1992)337
Plzák, Z., see Grüner, B. 595(1992)169
Reiss, W. H., see Jones, L. A. 595(1992)209
Reiss, W. H., see Jones, L. A. 595(1992)221
Reya, R., see Hlavačková, E. 595(1992)137
Richardson, D. E., Bremner, J. B. and O'Grady, B. V.
Quantitative analysis of total resin acids by high-performance liquid chromatography of their coumarin ester derivatives 595(1992)155
Roth, M., see Koliadima, A. 595(1992)237
Rühl, C., see Jiao, Y. 595(1992)364
Sadlej-Sosnowska, N. and Śledzińska, I.
Validation of chromatographic retention models in reversed-phase high-performance liquid chromatography by fitting experimental data to the relevant equations 595(1992)53
Saito, K., Horie, M., Nose, N., Nakagomi, K. and Nakazawa, H.
High-performance liquid chromatography of histamine and 1-methylhistamine with on-column fluorescence derivatization 595(1992)163
Sander, L. C., see Epler, K. S. 595(1992)89
Sapis, J.-C., see Voirin, S. G. 595(1992)269
Schwadorf, K. and Müller, H.-M.
Determination of α- and β-zearalenol and zearalenone in cereals by gas chromatography with ion-trap detection 595(1992)259
Schwer, C., see Kenndler, E. 595(1992)313
Shigenobu, K., see Koizumi, K. 595(1992)340
Shin, M. J., see Park, J. H. 595(1992)45
Śledzińska, I., see Sadlej-Sosnowska, N. 595(1992)53
Tadano, J., see Kawasaki, S. 595(1992)193
Takeuchi, T. and Nagae, N.
Stationary phase effect on enantioselectivity of dansyl phenylalanine in microcolumn liquid chromatography with γ-cyclodextrin as mobile phase additive 595(1992)121
Tanaka, N., see Kimata, K. 595(1992)77
Tanimoto, T., see Koizumi, K. 595(1992)340
Tharakan, J., Highsmith, F., Clark, D. and Drohan, W.
Physical and biochemical characterization of five commercial resins for immunoaffinity purification of factor IX 595(1992)103
Traag, W. A., see Van Rhijn, J. A. 595(1992)289
Tuinstra, L. G. M. T., see Van Rhijn, J. A. 595(1992)289
Ueda, H., see Kawasaki, S. 595(1992)193

Van Rhijn, J. A., Traag, W. A., Kulik, W. and Tuinstra, L. G. M. T.
Automated clean-up procedure for the gas chromatographic–high-resolution mass spectrometric determination of polychlorinated dibenzo-*p*-dioxins and dibenzofurans in milk 595(1992)289
Vinš, I., see Grüner, B. 595(1992)169
Voirin, S. G., Baumes, R. L., Sapis, J.-C. and Bayonove, C. L.
Analytical methods for monoterpene glycosides in grape and wine. II. Qualitative and quantitative determination of monoterpene glycosides in grape 595(1992)269
Waisser, K., see Klimeš, J. 595(1992)334
Walker, L. V., Walsh, J. R. and Webber, J. J.
High-performance liquid chromatography of sulphonamides extracted from bovine and porcine muscle by solid-phase dispersion 595(1992)179
Walsh, J. R., see Walker, L. V. 595(1992)179
Webber, J. J., see Walker, L. V. 595(1992)179
Weber, R., see Jiao, Y. 595(1992)364
Wise, S. A., see Epler, K. S. 595(1992)89
Wolf, R. M., see Francotte, E. 595(1992)63
Wu, H.-L., see Chen, S.-H. 595(1992)203
Wu, S.-M., see Chen, S.-H. 595(1992)203
Yang, S. S.
Determination of maleic hydrazide in tobacco by micellar liquid chromatography 595(1992)346
Yeung, E. S., see Lee, T. T. 595(1992)319
Ziegler, R. G., see Epler, K. S. 595(1992)89

Errata

J. Chromatogr., 587 (1991) 338–342
Page 342, ref. 1 should read "P. Werkhoff and W. Bretschneider, *J. Chromatogr.*, 405 (1987) 87".

J. Chromatogr., 589 (1992) 31–43
Page 31, name of third author should read "H. Piryns".

J. Chromatogr., 590 (1992) 263–270
Page 264, *Standard metal solutions* section, 5th line: "(II) chloride" should read "managanese(II) chloride".